
United States Department of Agriculture

Natural Resources Conservation Service

East Central Glaciated Regional MLRA Soil Survey Office Indianapolis, IN

Classification and Correlation of Soils in Owen County, Indiana



OWEN COUNTY, INDIANA
United States Department of Agriculture
Natural Resources Conservation Service

Classification and Correlation of the Soils of Owen County, Indiana

A Subset of MLRA's 114, 120 and 122

This correlation was prepared by Byron G. Nagel MLRA Project Leader, in February 1997. A preliminary correlation was conducted August 29 - September 2, 1994, at the Soil Conservation Field Office in Spencer, Indiana. Full time participants were Byron G. Nagel, Assistant State Soil Scientist, SCS; and Paul McCarter Jr., Project Leader, SCS. Other participants were project team members, Jerold Shively, Area Soil Scientist, SCS; and Neil Patterson, Soil Scientist, SCS. Carl Glocker, Soil Scientist, SSQA Staff, SCS, NSSC participated in the Final Field Review held October 4-8, 1993. Gary Struben, Soil Data Quality Specialist, MLRA Region 11, Indianapolis, Indiana made the technical review of this document.

In preparing this correlation, the following was available: 1) soil survey text manuscript, 2) soil maps, 3) field notes and transect data, 4) soil correlation samples, 5) laboratory data, 6) soil interpretation records, and 7) SOI-6 file.

Headnote for Detailed Soil Survey Legend

Map symbols consist of a combination of letters, or letters and numbers. The first capital letter is the initial one of the map unit name. Then two lower case letters that follow separate the map units having names that begin with the same letter. The second capital letter indicates the slope class. Symbols without a slope letter are for miscellaneous areas. Symbols ending with a number indicate the erosion class. Symbols ending with a capital letter as the fifth character indicates inundation phases or other soil phases.

SOIL CORRELATION OF OWEN COUNTY, INDIANA

Field symbols	Field map unit name	Publi- cation symbol	unit name
AdA, Pa, Ad	Adrian muck, warm, undrained, 0 to 1 percent slopes	AbgAU	Adrian muck, undrained, 0 to 1 percent slopes
AoC, ByC	Alvin-Bloomfield loamy sands, 6 to 12 percent slopes	AfeC	Alvin-Bloomfield loamy sands, 6 to 12 percent slopes
AoD, ByD	Alvin-Bloomfield loamy sands, 12 to 18 percent slopes	AfeD	Alvin-Bloomfield loamy sands, 12 to 18 percent slopes
AOE, ByE, AnE, PrE2, ByF, ByG, AOF	Alvin-Bloomfield- Princeton complex, 18 to 25 percent slopes	AffE	Alvin-Bloomfield- Princeton complex, 18 to 25 percent slopes
AnB, PrB, AoB	Alvin-Princeton fine sandy loams, 2 to 6 percent slopes	AfsB	Alvin-Princeton fine sandy loams, 2 to 6 percent slopes
AnC, PrC, AnC2	Alvin-Princeton fine sandy loams, 6 to 12 percent slopes	,	Alvin-Princeton fine sandy loams, 6 to 12 percent slopes
AnD, PrD2, AnD2	Alvin-Princeton fine sandy loams, 12 to 18 percent slopes	AftD	Alvin-Princeton sandy loams, 12 to 18 percent slopes
ArA, Ar	Armiesburg silt loam, 0 to 2 percent slopes, occasionally flooded, brief duration	į	Armiesburg silty clay loam, 0 to 2 percent slopes, occasionally flood- ed, brief duration
MuB2, MuA, MuB	Muren Variant silt loam, 2 to 6 percent slopes, eroded		Arney silt loam, 2 to 6 percent slopes, eroded
AvB2, AwB3, AxB2, AvB AxB	Ava silt loam, 2 to 6 percent slopes, eroded	AloB2	Ava silt loam, 2 to 6 percent slopes, eroded
AyA, RoA, Wt, Ay	Ayrshire fine sandy loam, 0 to 2 percent slopes	AmkA	Ayrshire fine sandy loam, 0 to 2 percent slopes

Field symbols	Field map unit name	Publi- cation symbol	unit name
SoA, So	Stendal silt loam, 0 to 2 percent slopes, frequently flooded, very brief duration		Belknap silt loam, 0 to 2 percent slopes, frequently flooded, brief duration
BgA, Bg	Birds silt loam, 0 to 1 percent slopes, frequently flooded, very brief duration	BgeAH	Birds silt loam, 0 to 1 percent slopes, frequently flooded, brief duration
BzA, Bz	Bonnie silt loam, 0 to 1 percent slopes, frequently flooded, brief duration	BodAH	Bonnie silt loam, 0 to 1 percent slopes, frequently flooded, very brief duration
CoB2, CsB2, CrB2, CcB2, CoB, CbB2	Cincinnati silt loam, 2 to 6 percent slopes, eroded	CkkB2	Cincinnati silt loam, 2 to 6 percent slopes, eroded
CoC2, CpC2	Cincinnati silt loam, 6 to 12 percent slopes, eroded	CkkC2	Cincinnati silt loam, 6 to 12 percent slopes, eroded
CpC3, CoC3, CtC3, CfC3	Cincinnati silt loam, 6 to 12 percent slopes, severely eroded	CkkC3	Cincinnati silt loam, 6 to 12 percent slopes, severely eroded
CoD2, HcD2, CtD2, CpD2, HkD2, ChD2	Cincinnati silt loam, 12 to 18 percent slopes, eroded	CkkD2	Cincinnati silt loam, 12 to 18 percent slopes, eroded
CpD3, HkD3, CfD3, HdD3	Cincinnati silt loam, 12 to 18 percent slopes, severely eroded	CkkD3	Cincinnati silt loam, 12 to 18 percent slopes, severely eroded
CvC2, CvB2	Crider silt loam, 6 to 12 percent slopes, eroded	CspC2	Crider silt loam, 6 to 12 percent slopes, eroded

Field symbols	Field map unit name	Publi- cation symbol	unit name
CwC3, CvC3	Crider silt loam, 6 to 12 percent slopes, severely eroded	CspC3	Crider silt loam, 6 to 12 percent slopes, severely eroded
DbA, BaA, DcA, Ba	Dubois silt loam, 0 to 2 percent slopes	DfnA	Dubois silt loam, 0 to 2 percent slopes
EbD2, EbC2	Ebal silt loam, 12 to 18 percent slopes, eroded	EaaD2	Ebal silt loam, 12 to 18 percent slopes, eroded
CkE2, CmE3, ChE, ChE2, CmE2, CkE, PbE2	Chetwynd Variant loam, 18 to 25 percent slopes, eroded	GaaE2	Gallimore loam, 18 to 25 percent slopes, eroded
ChG, ChF, CkF, CkG	Chetwynd Variant- Chetwynd loams, 25 to 70 percent slopes	GabG	Gallimore-Chetwynd loams, 25 to 70 percent slopes
JsA, Js, Gg, Gm	Gessie silt loam, 0 to 2 percent slopes, frequently flooded, brief duration	GblAH	Gessie silt loam, 0 to 2 percent slopes, frequently flooded, brief duration
GsD3, RyD3, GrD3, RxD3	Grayford silt loam, 12 to 18 percent slopes, severely eroded	GmcD3	Grayford silt loam, 12 to 18 percent slopes, severely eroded
GrE2, GrE, GrF, AlE, GsE3	Grayford silt loam, 18 to 25 percent slopes, eroded	GmcE2	Grayford silt loam, 18 to 25 percent slopes, eroded
GrD2, RxD2, AlD2	Grayford-Ryker silt loams, 12 to 18 percent slopes, eroded	GmhD2	Grayford-Ryker silt loams, 12 to 18 percent slopes, eroded
OmE2, OnE2 OwE3, OmE, OwE3, OnE,	Otwell Variant silt loam, 18 to 25 percent slopes, eroded	GmpE2	Greybrook silt loam, 18 to 25 percent slopes, eroded

Field map unit name	Publi- cation symbol	unit name
Otwell Variant silt loam, 25 to 35 percent slopes	GmpF	Greybrook silt loam, 25 to 35 percer slopes
Hagerstown silt loam, 12 to 18 percent slopes, eroded	HarD2	Haggatt silt loam, 12 to 18 percent slopes, eroded
Hagerstown silt loam, 12 to 18 percent slopes, severely eroded	HarD3	Haggatt silt loam, 12 to 18 percent slopes, severely eroded
		Haggatt-Caneyville silt loams, 18 to 25 percent slopes, eroded
Haubstadt silt loam, 0 to 2 percent slopes	HCCA	Haubstadt silt loam, 0 to 2 percent slopes
Haubstadt silt loam, 2 to 6 percent slopes, eroded	НссВ2	Haubstadt silt loam, 2 to 6 percent slopes, eroded
Haubstadt silt loam, 6 to 12 percent slopes, severely eroded	HccC3	Haubstadt silt loam, 6 to 12 percent slopes, severely eroded
Haymond silt loam, 0 to 2 percent slopes, frequently flooded, brief duration	 HcgAH 	Haymond silt loam, 0 to 2 percent slopes frequently flooded, brief duration
Haymond silt loam, sinkhole, 0 to 2 percent slopes, occasionally ponded, very brief duration	HchA	Haymond silt loam, sinkhole, 0 to 2 percent slopes
	Otwell Variant silt loam, 25 to 35 percent slopes Hagerstown silt loam, 12 to 18 percent slopes, eroded Hagerstown silt loam, 12 to 18 percent slopes, severely eroded Hagerstown-Caneyville silt loams, 18 to 25 percent slopes, eroded Haubstadt silt loam, 0 to 2 percent slopes Haubstadt silt loam, 2 to 6 percent slopes, eroded Haubstadt silt loam, 6 to 12 percent slopes, eroded Haubstadt silt loam, 6 to 12 percent slopes, severely eroded Haymond silt loam, 6 to 2 percent slopes, frequently flooded, brief duration Haymond silt loam, sinkhole, 0 to 2 percent slopes, occasionally ponded,	Otwell Variant silt loam, 25 to 35 percent slopes Hagerstown silt loam, 12 to 18 percent slopes, eroded Hagerstown silt loam, 12 to 18 percent slopes, severely eroded Hagerstown-Caneyville silt loams, 18 to 25 percent slopes, eroded Haubstadt silt loam, 0 to 2 percent slopes, severely eroded Haubstadt silt loam, HccB2 Haubstadt silt loam, HccB2 Haubstadt silt loam, HccB2 The slopes aroded Haubstadt silt loam, HccB2 Haubstadt silt loam, HccB2 The slopes aroded Haubstadt silt loam, HccB2 Haubstadt silt loam, HccB3 Haubstadt silt loam, HccB3 Haubstadt silt loam, HccB3 Haubstadt silt loam, HccB3 Hauppers aroded Haymond silt loam, HccB3 Haymond silt loam, HcgAH The slopes, frequently flooded, brief duration Haymond silt loam, Sinkhole, O to 2 The percent slopes, occasionally ponded,

OWEN COUNTY, INDIANA --Continued

Field symbols	Field map unit name	Publi- cation symbol	unit name
HeG	Hickory loam, 45 to 70 percent slopes	HefG	Hickory loam, 45 to 70 percent slopes
HrE2, HrE, HrD2, HrD3	Hickory-Ryker Variant silt loams, karst, hilly, eroded	HelD2	Hickory-Stinesville silt loams, karst, hilly, eroded
HcE, HcE2, HkE2, CoE2, HkE, HeE2, HdE	Hickory silt loam, 18 to 25 percent slopes		Hickory silt loam, 18 to 25 percent slopes
HkE3, HpE3	Hickory silt loam, 18 to 25 percent slopes, severely eroded	HeoE3	Hickory silt loam, 18 to 25 percent slopes, severely eroded
HcF, HcG, Hef	Hickory silt loam, 25 to 60 percent slopes		Hickory silt loam, 25 to 70 percent slopes
HmG, HmF, HnG	Hickory-Gilpin complex, 35 to 60 percent slopes	HepG	Hickory-Adyeville complex, 35 to 60 percent slopes
HdG, HdF, HgG	Hickory-Chetwynd loams, 35 to 70 percent slopes	HesG	Hickory-Chetwynd loams, 35 to 70 percent slopes
HnE, HpE2, HnE2	Hickory-Rosine Variant silt loams, 18 to 25 percent slopes	HeuE	Hickory-Wellston silt loams, 18 to 25 percent slopes
HnF	Hickory-Rosine Variant silt loams, 25 to 35 percent slopes	HeuF	Hickory-Wellston silt loams, 25 to 35 percent slopes
FeB	Fairpoint Variant silt loam, 1 to 5 percent slopes	HkaB	Hollybrook silt loam, 1 to 5 percent slopes
HsA, Ht, Hs, WsA	Holton silt loam, 0 to 2 percent slopes, frequently flooded, very brief duration		Holton silt loam, 0 to 2 percent slopes frequently flooded, very brief duration

	Field symbols	Field map unit name	Publi- cation symbol	unit name
HuA,	Hu	Hoosierville silt loam, 0 to 1 percent slopes	HnoA	Hoosierville silt loam, 0 to 1 percent slopes
IvA		Iva Variant silt loam, 0 to 2 percent slopes	LnfA	Lieber silt loam, 0 to 2 percent slopes
IvB,	IvB2	Iva Variant silt loam, 2 to 4 percent slopes	LnfB	Lieber silt loam, 2 to 4 percent slopes
McA		Martinsville loam, 0 to 2 percent slopes, rarely flooded	MecAQ	Martinsville loam, 0 to 2 percent slopes, rarely flooded
MeA,	MfA	Martinsville loam, 0 to 2 percent slopes	MecA	Martinsville loam, 0 to 2 percent slopes
MeB,	MeB2	Martinsville loam, 2 to 6 percent slopes	MecB	Martinsville loam, 2 to 6 percent slopes
HfA,	Нf	Nolin Variant silt loam, 0 to 2 percent slopes, frequently flooded, brief duration	!	McAdoo silt loam, 0 to 2 percent slopes, frequently flooded, brief duration
NxA,	Nx	Nolin Variant silt loam, 0 to 2 percent slopes, occasionally flooded, brief duration	İ	McAdoo silt loam, 0 to 2 percent slopes occasionally flood- ed, brief duration
MgA		McGary silt loam, 0 to 2 percent slopes	MhuA	McGary silt loam, 0 to 2 percent slopes
FcG,	FgG	Fairpoint Variant silty clay loam, 35 to 75 percent slopes	MrcG	Minnehaha silty clay loam, 35 to 75 percent slopes
MhA,	Mh	Montgomery silty clay loam, 0 to 1 percent slopes		Montgomery silty clay loam, 0 to 1 percent slopes

Field symbols	Field map unit name	 Publi- cation symbol	unit name
MnA, Mn	Moundhaven loamy fine sand, 0 to 2 percent slopes, frequently flooded, brief duration		Moundhaven loamy fine sand, 0 to 2 percent slopes, frequently flooded, brief duration
NaB	Nawakwa, silt loam, 2 to 8 percent slopes	NaeB	Nawakwa silt loam, 2 to 8 percent slopes
NeA, WaA, Wa, Sm, Wb, WbA, Sh, Ne	Newark silty clay loam, 0 to 2 percent slopes, frequently flooded, brief duration	,	Newark silt loam, 0 to 2 percent slopes, frequently flooded, brief duration
OfA, Of, Ph, PhA, Oh	Oldenburg silt loam, 0 to 2 percent slopes, frequently flooded, very brief duration	OfaAH	Oldenburg silt loam, 0 to 2 percent slopes, frequently flooded, very brief duration
OgA, Og	Oldenburg fine sandy loam, 0 to 2 percent slopes, sandy substratum, frequently flooded, very brief duration		Oldenburg fine sandy lo 0 to 2 percent slopes, sandy substratum, frequently flooded, ve brief duration
EmB, EmA, EmB2, EnB	Elkinsville Variant silt loam, 1 to 3 percent slopes	OfeB	Olephant silt loam, 1 to 3 percent slopes
OmC2, OtC2	Otwell silt loam, 6 to 12 percent slopes, eroded	OmkC2	Otwell silt loam, 6 to 12 percent slopes, eroded
OwC3, OmC3, OtC3, OpC3, OnC3	Otwell silt loam, 6 to 12 percent slopes, severely eroded	OmkC3	Otwell silt loam, 6 to 12 percent slopes, severely eroded
OwD3, OmD2, OmD3, OwD2, OnD2	Otwell silt loam, 12 to 18 percent slopes, severely eroded	OmkD3	Otwell silt loam, 12 to 18 percent slopes, severely eroded

Field symbols	Field map unit name	Publi- cation symbol	unit name
PbC2, PkC2, PaC2, PcC PcC2, PbC,	Parke silt loam, 6 to 12 percent slopes, eroded	PbbC2	Parke silt loam, 6 to 12 percent slopes, eroded
PcC3, PbC3, CkC3	Parke silt loam, 6 to 12 percent slopes, severely eroded	PbbC3	Parke silt loam, 6 to 12 percent slopes severely eroded
PbD2, PkD2, PcD2, PaD2	Parke silt loam, 12 to 18 percent slopes, eroded	PbbD2	Parke silt loam, 12 to 18 percent slopes, eroded
PcD3, PbD3	Parke silt loam, 12 to 18 percent slopes, severely eroded	PbbD3	Parke silt loam, 12 to 18 percent slopes, severely eroded
JoA	Johnsburg Variant silt loam, 0 to 2 percent slopes	PcnA	Patricksburg silt loam, 0 to 2 percent slopes
BbA	Pekin silt loam, 0 to 2 percent slopes, rarely flooded, very brief duration	j	Pekin silt loam, 0 to 2 percent slopes, rarely flooded
PsB2, PdB2	Pekin silt loam, 2 to 6 percent slopes, eroded	PcrB2	Pekin silt loam, 2 to 6 percent slopes, eroded
PsC2	Pekin silt loam, 6 to 12 percent slopes, eroded	PcrC2	Pekin silt loam, 6 to 12 percent slopes, eroded
PeA, Vn, PgA, Ra, ReA, Pe	Peoga silt loam, 0 to 1 percent slopes	PhaA	Peoga silt loam, 0 to 1 percent slopes
PnA	Peoga silt loam, ponded, 0 to 1 percent slopes	PhaAP	Peoga silt loam, ponded, 0 to 1 percent slopes

OWEN COUNTY, INDIANA --Continued

Field symbols	Field map unit name	Publi- cation symbol	unit name
PmA, Pm	Piankeshaw silt loam, 0 to 2 percent slopes, frequently flooded, very brief duration	PlcAH	Piankeshaw silt loam 0 to 2 percent slopes, frequently flooded, very brief duration
PkA, PbA	Pike silt loam, 0 to 2 percent slopes	PlfA	Pike silt loam, 0 to 2 percent slopes
PkB2, PbB2, PbB, PaB2, PkB, CkB2	 Pike silt loam, 2 to 6 percent slopes, eroded	PlfB2	Pike silt loam, 2 to 6 percent slopes, eroded
Рр	Pits, quarry	Pml	Pits, quarries
PoA, Po	Pope fine sandy loam, 0 to 2 percent slopes, frequently flooded, very brief duration	PrwAH	Pope fine sandy loam 0 to 2 percent slopes, frequently flooded, very brief duration
TsB, TsB2, TsA	Tilsit Variant silt loam, 1 to 3 percent slopes, eroded	PryB	Potawatomi silt loam, 1 to 3 percent slopes
MaD3, MdD3, MaD2, MdD2	Markland Variant silt loam, 12 to 18 percent slopes, severely eroded	PsaD3	Pottersville silt loam, 12 to 18 percent slopes, severely eroded
MaF, MaE, MaE2, MdE2	Markland Variant, 25 to 50 percent slopes	PsaG	Pottersville silt loam, 25 to 50 percent slopes
EnF, EnE, OtE3	Markland Variant silt loam, karst, steep	PsbF	Pottersville silt loam, karst, steep
SaA, Sa	Riverwash	Rnn	Riverwash
CsG, GrG, CbG, CoG	Corydon Variant- Corydon-Rock outcrop complex, 35 to 60 percent slopes	RpzG	Romona-Corydon-Rock outcrop complex, 35 to 60 percent slopes
RyC3, RxC3, AlC3, GsC3, GrC3	Ryker silt loam, 6 to 12 percent slopes, severely eroded	RtcC3	Ryker silt loam, 6 to 12 percent slopes, severely eroded

Field symbols	Field map unit name	Publi- cation symbol	unit name
ScA, AvA	Shakamak silt loam, 1 to 3 percent slopes	SfoA	Shakamak silt loam, 1 to 3 percent slopes
MaB2	Shircliff silt loam, 2 to 6 percent slopes, eroded	sfyB2	Shircliff silt loam, 2 to 6 percent slopes, eroded
MaC2, MdC3, MdC2, MaC3, MdC	Shircliff silt loam, 6 to 12 percent slopes, eroded	SfyC2	Shircliff silt loam, 6 to 12 percent slopes, eroded
CsC2, CrC2	Cincinnati Variant silt loam, 6 to 12 percent slopes, eroded	SneC2	Solsberry silt loam, 6 to 12 percent slopes, eroded
CrC3, CsC3, HpC3	Cincinnati Variant silt loam, 6 to 12 percent slopes, severely eroded	SneC3	Solsberry silt loam, 6 to 12 percent slopes, severely eroded
CsD2, HnD2, CrD2	Cincinnati Variant silt loam, 12 to 18 percent slopes, eroded	SneD2	Solsberry silt loam, 12 to 18 percent slopes, eroded
CrD3, HpD3, CsD3, HnD3	Cincinnati Variant silt loam, 12 to 18 percent slopes, severely eroded	SneD3	Solsberry silt loam, 12 to 18 percent slopes, severely eroded
UbD4, Ub	Cincinnati Variant silt loam, gullied, 12 to 18 percent slopes	SneD5	Solsberry silt loam, 12 to 18 percent slopes, gullied
SnA, Sn	Steff silt loam, 0 to 2 percent slopes, frequently flooded, very brief duration	StaAH	Steff silt loam, 0 to 2 percent slopes, frequently flooded, very brief duration
StdAH	Stendal silt loam, 0 2 percent slopes, frequently flooded, brief duration	StdAH	Stendal silt loam, 0 to 2 percent slopes, frequently flooded, brief duration
RxB2, GrA, RxA, GrB2, AlB2, AlA, RyB2, GrB	Ryker Variant silt loam, 2 to 6 percent slopes, eroded	!	Stinesville silt loam, 2 to 6 percent slopes, eroded

OWEN COUNTY, INDIANA -- Continued

Field symbols	Field map unit name	Publi- cation symbol	unit name
RxC2, GrC2, AlC2, RyC2, RcC2	Ryker Variant silt loam, 6 to 12 percent slopes, eroded	StfC2	Stinesville silt loam, 6 to 12 percent slopes, eroded
RzC2, RzB2	Ryker Variant-Ryker- Grayford silt loams, karst, rolling, eroded	StgC2	Stinesville-Ryker- Grayford silt loams, karst, rolling, eroded
GtD2, GtE2, GtD3	Ryker Variant-Ryker- Grayford silt loams, karst, hilly, eroded	ĺ	Stinesville-Ryker- Grayford silt loams, karst, hilly, eroded
SpA, Sp	Stonelick sandy loam, 0 to 2 percent slopes, frequently flooded, brief duration	SupAH	Stonelick sandy loam, 0 to 2 percent slopes, frequently flooded, brief duration
HhG, OtG, OtF, OfG, HhF	Uniontown Variant- Hickory silt loams, 35 to 70 percent slopes	SwhG	Stubenville-Hickory complex, 35 to 70 percent slopes
TaA, Ta	Taggart silt loam, 0 to 2 percent slopes	TadA	Taggart silt loam, 0 to 2 percent slopes
FgB, FgA	Fairpoint Variant silt loam, 2 to 8 percent slopes	TakB	Tapawingo silt loam, 2 to 8 percent slopes
BfG, WeG	Berks-Rock outcrop complex, 35 to 70 percent slopes	TcgG	Tipsaw-Rock outcrop complex, 35 to 70 percent slopes
BeF, BeG	Rosine Variant-Tipsaw complex, 25 to 60 percent slopes	TtaG	Tulip-Tipsaw complex, 25 to 60 percent slopes
WpE, WpE2, EbE, EbE2, WmE, WmE2, WnE, WnE2 WnE3, WpE3	Rosine Variant- Wellston-Gilpin complex, 18 to 25 percent slopes	TtcE	Tulip-Wellston- Adyeville silt loams, 18 to 25 percent slopes

Field symbols	Field map unit name	Publi- cation symbol	unit name
,,	Udorthents, 0 to 2 percent slopes	Uaa	Udorthents, cut and filled
VgA	Vigo silt loam, 0 to 2 percent slopes	VdgA	Vigo silt loam, 0 to 2 percent slopes
W, Wct	Water	W	Water
WdA, OcA, OkA, EvA, Nv	Wea, sandy substratum-Elston complex, 0 to 2 percent slopes	WeaA	Wea, sandy substratum-Elston complex, 0 to 2 percent slopes
WfA, ObB2, Ob	Wea loam, sandy substratum-Elston complex, 0 to 2 percent slopes, rarely flooded	WeaAQ	Wea, sandy substratum-Elston complex, 0 to 2 percent slopes, rarely flooded
WmD2, WnD2	Wellston silt loam, 12 to 18 percent slopes, eroded	WhfD2	Wellston silt loam, 12 to 18 percent slopes, eroded
WtA	Whitaker loam, 0 to 2 percent slopes, occasionally flooded, brief duration	WlmAK	Whitaker loam, 0 to 2 percent slopes, occasionally flooded, brief duration
WhA, Wm, Wk, WkA, Es, Lo, Wh	Wilbur silt loam, 0 to 2 percent slopes, frequently flooded, brief duration	•	Wilbur silt loam, 0 to 2 percent slopes, frequently flooded, brief duration
WvA, Wv	Wilhite silty clay, 0 to 1 percent slopes, frequently flooded, long duration		Wilhite silty clay, 0 to 1 percent slopes frequently flooded, long duration
PfA, Pf	Wilhite silty clay loam, 0 to 1 percent slopes, frequently flooded, brief duration	:	Wilhite silty clay loam, 0 to 1 percent slopes, frequently flooded, brief duration

Field symbols	Field map unit name	Publi- cation symbol	unit name
WzA, Wz	Wirt loam, 0 to 2 percent slopes, rarely flooded, brief duration	WprAQ	Wirt loam, 0 to 2 percent slopes, rarely flooded
WxA, Wx, Cy, Cx, Cu	Wirt silt loam, 0 to 2 percent slopes, frequently flooded, very brief duration	WpuAH	Wirt silt loam, 0 to 2 percent slopes, frequently flooded, very brief duration
WyA, Wy, Cf	Wirt silt loam, 0 to 2 percent slopes, occasionally flooded, brief duration	WpuAK	Wirt silt loam, 0 to 2 percent slopes, occasionally flooded, brief duration
PdB, PdD2, PdD, WcA, UnB, Wc	Wirt-Pekin silt loams, 0 to 6 percent slopes, frequently flooded, very long duration	WpyBJ	Wirt-Pekin silt loams, 0 to 6 percent slopes, frequently flooded, very long duration
ZaB2, ZaB, ZnB3, ZnB2	Zanesville silt loam, 2 to 6 percent slopes, eroded	ZamB2	Zanesville silt loam, soft bedrock substratum, 2 to 6 percent slopes, eroded
ZaC2, WmC2, ZnC2	Zanesville silt loam, 6 to 12 percent slopes, eroded	ZamC2	Zanesville silt loam, soft bedrock substratum, 6 to 12 percent slopes, eroded
ZnC3, WnC3, ZaC3, ZmC3,	Zanesville silt loam, 6 to 12 percent slopes, severely eroded	ZamC3	Zanesville silt loam, soft bedrock substratum, 6 to 12 percent slopes, severely eroded

OWEN COUNTY, INDIANA --Continued

Field symbols	Field map unit name	Publi- cation symbol	unit name
ZaD2, ZnD2	Zanesville silt loam, 12 to 18 percent slopes, eroded	ZamD2	Zanesville silt loam, soft bedrock substratum, 12 to 18 percent slopes, eroded
UgD4, Ug	Zanesville silt loam, gullied, 12 to 18 percent slopes	ZamD5	Zanesville silt loam, soft bedrock substratum, 12 to 18 percent slopes, gullied
ZnD3, WnD3 WmD3, WeD3, WpD3, ZaD3,	Zanesville-Rosine Variant silt loams, 12 to 18 percent slopes, severely eroded	ZapD3	Zanesville, soft bedrock substratum- Tulip silt loams, 12 to 18 percent slopes, severely eroded

Series Established by this Correlation and County of Type Location

Adyeville (Perry Co.)

Patricksburg (Owen Co.)

Arney (Clay Co.)

Potawatomi (Owen Co.)

Gallimore (Owen Co.)

Pottersville (Owen Co.)

Greybrook (Owen Co.)

Romona (Owen Co.)

Haggatt (Washington Co.)

Solsberry (Owen Co.)

Hollybrook (Owen Co.)

Stinesville (Owen co.)

Lieber (Clay Co.)

Stubenville (Owen co.)

McAdoo (Posey Co.)

Tapawingo (Owen Co.)

Minnehaha (Owen Co.)

Tipsaw (Perry Co.)

Nawakawa (Owen Co.)

Tulip (Owen Co.)

Olephant (Owen Co.)

Series Dropped or Made Inactive

None

Cooperators' Names and Credits

The cooperators for the front cover are

United States Department of Agriculture Natural Resources Conservation Service in cooperation with the Purdue University Agricultural Experiment Station

The credits to be given on page ii of the published soil survey are as follows:

This survey was made cooperatively by the National Resources Conservation Service and the Purdue University Agricultural Experiment Station. It is part of the technical assistance furnished to the Owen County Soil and Water Conservation District.

Prior Soil Survey Publications

The last soil survey of Owen County was completed in 1959 and published by the United States Department of Agriculture, Soil Conservation Service in October 1964. Reference to the prior soil survey will be included in the literature citation of the manuscript. This survey replaces the October 1964 soil survey, and provides additional data, updated soil interpretations and 1:12,000 scale soil maps on an orthophotographic base.

Explanation of Map Unit Symbol Characters in the Indiana Statewide Legend

The first three characters are alpha characters which indicate the soil series and phases. The fourth character is an alpha character which represents the slope class. The fifth character is an alpha character or numeric character and represents one of the following in Owen County:

- 2 Moderate Erosion Class
- 3 Severe Erosion Class
- 5 Gullied Phase
- H Frequently flooded, very brief or brief duration
- I Frequently flooded, long duration
- J Frequently flooded, very long duration
- K Occasionally flooded, brief duration
- P Ponded
- Q Rarely flooded
- U Undrained

Instructions for Map Compilation, Map Finishing, and Digitizing

Map compilation has been completed and the soil maps are being digitized at the NRCS Indianapolis MLRA Project Office. Selected county roads will be numbered. The field symbol OtC3 on soil map field sheets 4, 5, 6, 10 and 11 is to be converted to field symbol HbC3. The field symbols So and SoA on designated areas of field sheets 60, 61, 67, and 68 are to be converted to publication symbol StdAH. Field symbols WnE3 and WpE3 are to be converted to published symbol TtcE and a severely eroded spot symbol placed within the map unit. The special symbol for sinkholes will not be placed in these map units which are designated as karst or sinkhole phases: HchAP, HelD2, PsbF, StgC2, StgD2. Intermittent ditches are to be used in small valleys and in farm fields where the surrounding natural streams are intermittent. Perennial ditches are to be used in large valleys. All special symbols were evaluated for times used and consistency after they were digitized.

Conventional and Special Symbols Legend

Only those symbols indicated on the NRCS-SOILS-37A (7/96) will be shown on the legend and placed on the soil maps. The definition of the Marsh sandy spot, and wet spot special symbols are not defined, in Owen County, as is stated in Part 647 (11/96) of the National Soil Survey Handbook. Perennial water also includes miscellaneous water in Owen County.

CONVENTIONAL AND SPECIAL

Soil Survey Area: Owen County State: IN

SYMBOLS LEGEND

9/94 DESCRIPTION DESCRIPTION SYMBOL SYMBOL DESCRIPTION SYMBOL SPECIAL SYMBOLS FOR SOIL SURVEY CULTURAL FEATURES CULTURAL FEATURES (cont.) AND SSURGO SOIL DELINEATIONS AND SOIL SYMBOLS BOUNDARIES MISCELLANEOUS CULTURAL FEATURES National, state, or province Farmstead, house (amit in urban areas) T LANGFORM FEATURES Church ESCARPMENTS School Bedrock **74748474844747474747474747474747** Minor civil division Other than bedrock Other Religion (lobel) SHORT STEEP SLOPE ⊙ States GULLY ~~~~~~~~~ Reservation (national forest or park, Located object (lobel) DEPRESSION, closed Tank (label) SINKHOLE Leckout Tower Limit of soil survey (label) and/or denied access areas QII and/or Natural Gas Wells # EXCAVATIONS PITS g / Fleid sheet matchline & neatting Borrew pli Lighthouse Gravel pH Previously Published Survey Mine or quarry OTHER BOUNDARY (lobel) 1 00mg | 1 1 ± 1 Airport, airfield HYDROGRAPHIC FEATURES LANDERL Cemetery [E-20] [T] City/county park 1 Porenniol, double line I MISCELLANEOUS SURFACE FEATURES Control Blowout Perennial, single line STATE COORDINATE TICK Intermittent Gravelly spat ND DIVISION CORNERS Lova flow Drainage and GEOGRAPHIC COORDINATE TICK Marsh or swamp Rack autorop (includes sandsione and shale) TRANSPORTATION DRAINAGE AND IRRIGATION Soline spei 1 Double line congl (ighel) Divided roads Sandy spot Severely eroded spot Silde or ello Sodic spot Sooll erec SMALL LAKES, PONDS AND RESERVOIRS Stony spot ROAD EMBLEMS & DESIGNATIONS Yery stany spat Patenniol water Wei spat Interviate Very severely eroded Flood gool line Federal 010 spot T RECOMMENDED AD HOC SOIL SYMBOLS **√** State County, farm or ranch 374 п RAILROAD ٥ • POWER TRANSMISSION LINE (normally not shown) MISCELLANEOUS WATER FEATURES PIPELINE (normally not shown) FENCE (normally not shown) LEVEES Well, ortesion ٥ Without road management and the second † Single side slope (showing actual feature location) "V Prominent Hill or Peak Self Sample Site

18

DEFINITIONS OF SPECIAL FEATURES FOR OWEN COUNTY, INDIANA SOIL SURVEY

Feature	Label	Feature Definition
Escarpment, bedrock	ESB	A relatively continuous and steep, slope or cliff produced by erosion, or faulting, breaking the general continuity of more gently sloping land surfaces. Exposed material is hard or soft bedrock. Typically .5 to 2.5 acres.
Escarpment, other	ESO	A relatively continuous and steep, slope or cliff generally produced by erosion, but can be produced by faulting, breaking the general continuity of more gently sloping land surfaces. Exposed nonbedrock material is nonsoil or very shallow, poorly developed soil. Typically .5 to 2.5 acres.
Gravel pit	GPI	An open excavation from which soil and underlying material have been removed, and and used without crushing as a source of sand or of sand or gravel. Typically .5 to 2.5 acres.
Marsh	MAR	A small natural area, intermittently or permanently water-covered, and dominantly has hydrophytic vegetation. Typically .5 to 2.5 acres.
Mine or quarry	MPI	An open excavation from which soil and underlying material is removed exposing the bedrock. Typically 1 to 3 acres.
Perennial water	WAT	Small natural or manmade lake, pond, or pit that contains water most of the year. Typically .2 to 2.5 acres.
Rock outcrop	ROC	An exposure of bedrock at the surface of the earth. Not used where the named soils of the surrounding map unit are shallow over bedrock. Typically .1 to .2 acres.
Sandy spot	SAN	An area of soil where the surface layer is sandy loam, or loamy sand in an area where the surrounding soils have a silt loam or more clayey surface texture. Excluded are areas where the textural classes are adjoining, such as an area of loamy sand in a surrounding Map unit. Typically .25 to 2 acres.
Severely eroded spot	ERO	An area where on the average 75 percent or more of the original surface layer has been lost from accelerated erosion. Typically .5 to 2.5 acres.
Short, steep slope	SLP	Narrow soil area that has slopes that are at least 2 slope classes steeper than the slope class of the surrounding map units. Typically 1 to 2.5 acres.

Sinkhole

SNK A closed depression formed either by solution of the surficial rock or by collapse of underlying caves. Typically .5 to 2 acres.

Wet spot

WET An area of soil that is poorly drained or very poorly drained and that is at least 2 drainage classes wetter than the named soils in the surrounding map unit. Typically .5 to 1.75 acres.

Very severely eroded spot

An area where class 4 erosion exists. The original A and E and the upper B horizons are lost. Most areas consists of an intricate pattern of U-shaped gullies. The original soil can only be identified in areas adjacent to these very severely eroded spots. Typically .5 to 2 acres in size.

General Soil Map Units

The following 10 map units will be used on the general soil map legend:

McAdoo - Gessie - Stonelick Peoga - Dubois - Haubstadt Gallimore - Pike - Parke Ava - Solsberry - Hickory Zanesville - Tulip - Wellston Stinesville - Grayford Ava - Shakamak - Hickory Crider - Haggatt Wea - Elston - Martinsville Wirt - Holton - Stendal

CONVERSION LEGEND FOR OWEN COUNTY, INDIANA

Field symbol	Publi- cation symbol	Field symbol	Publi- cation symbol	Field symbol	Publi- cation symbol	Field symbol	Publi- cation symbol
Ad AdA Ala AlB2 AlC2	AbgAU AbgAU StfB2 StfB2 StfC2	ByD ByE ByF ByG Bz	AfeD AffE AffE AffE BodAH	CrC3 CrD2 CrD3 CsB2 CsC2	SneC3 SneD2 SneD3 CkkB2 SneC2	EvA FcG FeB FgA FgB	WeaA MrcG HkaB TakB TakB
AlC3 AlD2 AlE AnB AnC	RtcC3 GmhD2 GmcE2 AfsB AfsC	BzA CbB2 CbE CbE2 CbG	BodAH CkkB2 HasE2 HasE2 RpzG	CsC3 CsD2 CsD3 CsG CtC3	SneC3 SneD2 SneD3 RpzG CkkC3	FgG Gg Gm Gp GrA	MrcG GblAH GblAH Uaa StfB2
AnC2 AnD AnD2 AnE AoB	AfsC AftD AftD AffE AfsB	CcB2 Cf CfC3 CfD3 ChD2	CkkB2 WpuAK CkkC3 CkkD3 CkkD2	CtD2 Cu CvB2 CvC2 CvC3	CkkD2 WpuAH CspC2 CspC2 CspC3	GrB GrB2 GrC2 GrC3 GrD2	StfB2 StfB2 StfC2 RtcC3 GmhD2
AoC AoD AoE AoF Ar	AfeC AfeD AffE AffE AhrAK	ChE ChE2 ChF ChG CkB2	GaaE2 GaaE2 GabG GabG PlfB2	CVD2 CVE CVE2 CWC3 CWD2	HarD2 HasE2 HasE2 CspC3 HarD2	GrD3 GrE GrE2 GrF GrG	GmcD3 GmcE2 GmcE2 GmcE2 RpzG
ArA AvA AvB AvB2 AwB3	Ahrak Sfoa AloB2 AloB2 AloB2	CkC3 CkE CkE2 CkF	PbbC3 GaaE2 GaaE2 GabG GabG	CwD3 Cx Cy DbA DbB2	HarD3 WpuAH WpuAH DfnA HccB2	GsC3 GsD3 GsE3 GtD2 GtD3	RtcC3 GmcD3 GmcE2 StgD2 StgD2
AxB AxB2 Ay AyA Ba	AloB2 AloB2 AmkA AmkA DfnA	CmE2 CmE3 CoB CoB2 CoC2	GaaE2 GaaE2 CkkB2 CkkB2 CkkC2	DcA DcB2 EbC2 EbD2 EbE	DfnA HccB2 EaaD2 EaaD2 TtcE	GtE2 HaA HaB HaB2 HbA	StgD2 HccA HccB2 HccB2
BaA BbA BeF BeG BfG	DfnA PcrAQ TtaG TtaG TcgG	CoC3 CoD2 CoE2 CoG	CkkC3 CkkD2 HeoE RpzG CkkC2	EbE2 EmA EmB EmB2 EnB	TtcE OfeB OfeB OfeB OfeB	HbB HbB2 HbB3 HbC3 HcD2	HccB2 HccB2 HccB2 HccC3 CkkD2
Bg BgA ByC	BgeAH BgeAH AfeC	CpC3 CpD2 CpD3 CrB2 CrC2	CkkC3 CkkD2 CkkD3 CkkB2 SneC2	EnE EnF Es	PsbF PsbF WokAH	HcE HcE2 HcF HcG HdD3	HeoE HeoE HeoG HeoG CkkD3

OWEN COUNTY, INDIANA -- Continued

Field symbol	Publi- cation symbol	Field symbol	Publi- cation symbol	Field symbol	Publi- cation symbol	Field symbol	Publi- cation symbol
HdE	HeoE	HuA	HnoA	NaB	NaeB	OwD2	OmkD3
HdF	HesG	HzD2	HarD2	Ne	NbhAH	OwD3	OmkD3
HdG	HesG	HzD3	HarD3	NeA	NbhAH	OwE3	GmpE2
HeE2	HeoE	HzE	HasE2	Nv	WeaA	OwE3	GmpE2
HeF	HeoG	HzE2	HasE2	N×	MhhAK	Pa	AbgAU
HeG	HefG	HzE3	HasE2	NxA	MhhAK	PaB2	PlfB2
Hf	MhhAH	IvA	LnfA	0b	WeaAQ	PaC2	PbbC2
HfA	MhhAH	IvB	LnfB	ObB2	WeaAQ	PaD2	PbbD2
Hg	HchA	IvB2	LnfB	OcA	WeaA	PbA	PlfA
HgA	HchA	JoA	PcnA	Of	OfaAH	PbB	PlfB2
HgG	HesG]] Js	GblAH	OfA	OfaAH	PbB2	PlfB2
HhF	SwhG	JsA	GblAH	OfG	SwhG	PbC	PbbC2
HhG	SwhG	Lo	WokAH	0g	OfcAH	PbC2	PbbC2
HkD2	CkkD2	MaB2	SfyB2	OgA	OfcAH	PbC3	PbbC3
HkD3	CkkD3	MaC2	sfyC2	Oh	OfaAH	PbD2	PbbD2
HkE	HeoE	MaC3	SfyC2	OkA	WeaA	PbD3	PbbD3
HkE2	HeoE	MaD2	PsaD3	OmA	HccA	PbE2	GaaE2
HkE3	HeoE3	MaD3	PsaD3	OmB2	HccB2	PcC	PbbC2
HmF	HepG	MaE	PsaG	OmC2	OmkC2	PcC2	PbbC2
HmG	HepG	MaE2	PsaG	OmC3	OmkC3	PcC3	PbbC3
HnD2	SneD2	 MaF	PsaG	OmD2	OmkD3	l PcD2	PbbD2
HnD3	SneD3	McA	MecAQ	OmD3	OmkD3	PcD3	PbbD3
HnE	HeuE	MdC	SfyC2	OmE	GmpE2	PdB	WpyBJ
HnE2	HeuE	MdC2	sfyc2	OmE2	GmpE2	PdB2	PcrB2
HnF	HeuF	MdC3	SfyC2	ĺ			
		l		OmF	GmpF	PdD	WpyBc
HnG	HepG	MdD2	PsaD3	OmG	GmpF	PdD2	WpyBo
Но	HcgAH	MdD3	PsaD3	OnB2	HccB2	Pe	PhaA
HoA	HcgAH	MdE2	PsaG	OnC3	OmkC3	PeA	PhaA
HpC3	SneC3	MeA	MecA	OnD2	OmkD3	Pf	WomAl
HpD3	SneD3	MeB	MecB	OnE	GmpE2	PfA	WomAl
HpE2	HeuE	MeB2	MecB	OnE2	GmpE2	PgA	PhaA
нрег	HeoE3	MfA	MecA	OnF	GmpF	Ph	OfaAl
HrD2	HelD2	MgA	MhuA	OpC3	OmkC3	PhA	OfaAl
HrD3	HelD2	Mh	MsvA	OtC2	OmkC2	PkA	PlfA
HrE	HelD2	MhA	MsvA	İ		İ	
				OtC3	OmkC3	PkB	PlfB2
HrE2	HelD2	Mn	MvrAH	OtE3	PsbF	PkB2	PlfB
Hs	HleAH	MnA	MvrAH	OtF	SwhG	PkC2	PbbC:
HsA	HleAH	MuA	AkeB2	OtG	SwhG	PkD2	PbbD2
Ht	HleAH	MuB	AkeB2	OwC3	OmkC3	Pm	PlcA
Hu	HnoA	MuB2	AkeB2			!	

OWEN COUNTY, INDIANA --Continued

Field symbol	Publi- cation symbol	Field symbol	Publi- cation symbol	 Field symbol	Publi- cation symbol	Field symbol	Publi- cation symbol
PmA	PlcAH	TaA	TadA	WnC3	ZamC3		
PnA	PhaAP	TsA	PryB	WnD2	WhfD2		
Po	PrwAH	TsB	PryB	WnD3	ZapD3		
PoA	PrwAH	TsB2	PryB	WnE	TtcE		
Pp	Pm1	Ub	SneD5	WnE2	TtcE		
PrB	AfsB	UbD4	SneD5	WnE3	TtcE		
PrC	AfsC	ŬC	Uaa	WpD3	ZapD3	ĺ	
PrD2	AftD	UcA	Uaa	WpE	TtcE		
PrE2	AffE	Uđ	Uaa	WpE2	TtcE		
PsB2	PcrB2	UdA	Uaa	WpE3	TtcE		
D-00	D (10)	****	••-		**1 - * **		
PsC2	PcrC2	UdB	Uaa 	WsA	HleAH		
Ra	PhaA	UeA	Uaa	Wt	AmkA	Į	
RcC2	StfC2	Ug	ZamD5	WtA	WlmAK		
ReA	PhaA	UgD4	ZamD5	Wv	Wolai		
RoA	AmkA	UnB	WpyBJ	WvA	WolAI		
RxA	StfB2	VgA	VdgA	Wx	WpuAH	 	
RxB2	StfB2	Vn	PhaA	WxA	WpuAH		
RxC2	StfC2	i w	W	Wy	WpuAK	İ	
RxC3	RtcC3	Wa	NbhAH	WyA	WpuAK		
RxD2	GmhD2	WaA	NbhAH	Wz	WprAQ	Ì	
RxD3	GmcD3	 Wb	NbhAH	WzA	WprAQ		
	StfB2	WbA	NbhAH	1	ZamB2		
RyB2		1		ZaB			
RyC2	StfC2	WC	WpyBJ	ZaB2	ZamB2	!	
RyC3	RtcC3	Wct	W	ZaC2	ZamC2		
RyD3	GmcD3	WcA	WpyBJ				
RzB2	StgC2	WdA	WeaA	ZaD2	ZamD2		
RzC2	StgC2	WeD3	ZapD3	ZaD3	ZapD3	1	
Sa	Rnn	WeG	TcgG	ZmC3	ZamC3	1	
SaA	Rnn	WfA	WeaAQ	ZnB2	ZamB2	İ	
ScA	SfoA	Wh	WokAH	ZnB3	ZamB2	ĺ	
Sh	NbhAH	 WhA	WokAH	ZnC2	ZamC2		
Sm	NbhAH	Wk	WokAH	ZnC3	ZamC3	i	
Sn	StaAH	WkA	WokAH	ZnD2	ZamD2	i	
SnA	StaAH	Wm	WokAH	ZnD3	ZapD3	1	
5121	Dearm	WmC2	ZamC2		Lupbo		
So	BdxAH	WmD2	WhfD2				
		WmD3		1			
SoA	BdxAH		ZapD3	[
Sp	SupAH	WmE	TtcE				
SpA	SupAH	WmE2	TtcE	!			
StdAH	StdAH						
Ta	TadA					İ	

Classification of pedons sampled for Laboratory Analysis

Approved Series or	Sampled as	Soil Survey	Publication
Class Identification		Sample No.	Symbol
Alvin	Alvin	S91IN119-19	AfeC (c)
Alvin	Alvin	S91IN119-21	AfsC (c)
Armiesburg	Armiesburg	S90IN119-56	AhrAK (c)
Arney (a)	Muren	S93IN021-1	AkeB2 (c)
Ava	Ava	S87IN119-1	AloB2 (c)
Ava	Ava	S92IN119-4	AloB2 (c)
Belknap (d)	Stendal	S88IN119-35	BdxAH (c)
Bloomfield (d)	Bloomfield	S91IN119-20	AfeC (c)
Chetwynd	Chetwynd	S89IN119-15	GabG
Chetwynd (a)	Chetwynd	S93IN109-14	GabG (c)
Cincinnati	Cincinnati	S87IN119-3	CkkC3 (c)
Crider	Crider	S92IN119-1A	CspC2 (c)
Dubois	Dubois	S92IN119-1A	DfnA (c)
Ebal (d)	Ebal	S90IN119-55	EaaD2
Elston	Elston	S92IN119-33	
Gallimore (a)	Chetwynd Variant	S89IN119-16	WeaAQ (c) GabG
Gallimore (d)	Chetwynd Variant		
Gessie	Jules	S93IN109-13	GabG
Grayford (d)	Grayford	S88IN119-9	GblAH (c)
	Elkinsville Variant	S92IN119-26	GmcE2 (c)
Greybrook (a)		S91IN119-45	GmpF
Haggatt (d) Haubstadt	Hagerstown	S92IN119-25	HarD3
	Haubstadt	S92IN119-5	HccA
Hickory	Hickory	S87IN119-17	HeuF (c)
Hickory	Hickory	S92IN119-24	SwhG (c)
Hollybrook	Fairpoint Variant	S94IN119-2	HkaB
Hollybrook (a)	Fairpoint Variant	S94IN119-6	HkaB (c)
Hoosierville	Hoosierville	S92IN119-35	HnoA
Hoosierville (a)	Hoosierville	S93IN21-3	HnoA
Lieber	Iva Variant	S92IN119-36	LnfA
Lieber (a)	Iva Variant	S93IN21-2	LnfA
McAdoo (d)	Nolin	S90IN119-11	MhhAK (c)
McAdoo	Haymond	S92IN119-31	MhhAH (c)
Minnehaha (a)	Fairpoint Variant	S94IN119-5	MrcG (c)
Montgomery	Montgomery	S92IN119-53	MsvA
Nawakwa (a)	Fairpoint Variant	S94IN119-4	NaeB (c)
Newark	Wakeland	S91IN119-40	NbhAH (c)
Newark (b, d)	Newark	S87IN119-5	NbhAH
Oldenburg	Oldenburg	S90IN119-38	OfaAH (c)
Olephant (a)	Elkinsville Variant	S92IN119-22	OfeB
Parke (a)	Parke	S89IN119-12	PbbC2 (c)
Patricksburg (a)	Johnsburg Variant	S89IN119-8	PcnA
Pike	Pike	S89IN119-13	PlfA
Pike (a)	Pike	S92IN119-6	PlfA
Pike	Pike	S92IN119-7	PlfB2 (c)
Potawatomi	Tilsit Variant	S89IN119-6	PryB
Potawatomi (a)	Tilsit Variant	S92IN119-1	PryB (c)
Potawatomi (e)	Tilsit Variant	S92IN119-2	PryB
Pottersville (a)	Markland Variant	S92IN119-28	PsaG (c)

Classification of pedons sampled for Laboratory Analysis (continued)

Approved Series or	Sampled as	Soil Survey	Publication
Class Identification		Sample No.	Symbol
Pottersville (d) Princeton Romona (a) Shakamak (d)	Markland Variant	S90IN119-14	PsaD3
	Princeton	S91IN119-22	AfsC (c)
	Corydon Variant	S92IN119-32	RpzG (c)
	Shakamak	S87IN119-2	SfoA
Shircliff Solsberry (a) Stinesville	Markland	S90IN119-13	SfyB2 (c)
	Cincinnati Variant	S89IN119-14	SneC3 (c)
	Ryker Variant	S92IN119-8	StfB2 (c)
Stinesville (a, e) Stonelick Stonelick (d) Stubenville (a)	Ryker Variant Stonelick Stonelick Uniontown Variant	S92IN119-9 S91IN119-50 S89IN119-10 S92IN119-23	StfC2 (c) SupAH (c) SupAH SwhG (c)
Tapawingo (a) Tapawingo Tipsaw	Fairpoint Variant Fairpoint Variant Berks Variant	S94IN119-3 S94IN119-1 S90IN119-17	TakB (c) TakB TtaG (c)
Tulip (a) Tulip Vigo (a)	Rosine Variant Rosine Variant Vigo	S93IN119-34 S90IN119-46 S89IN119-11	TtaG (c) TtcE VdgA (c)
Vigo (e) Wea Wellston (d) Wellston (d)	Vigo	S92IN119-3	VdgA
	Wea	S92IN119-39	WeaAQ (c)
	Wellston	S90IN119-44	TtcE (c)
	Gilpin	S87IN119-18	HeuF (c)
Wilhite (d) Wilhite (b) Wirt	Wilhite Wilhite Wirt Wirt	S90IN119-1 S87IN119-4 S93IN119-1 S87IN119-1B	WolAI (c) WomAH (c) WpuAH (c)
Wirt (b)	Zanesville	S89IN119-1B	WpuAH
Zanesville		S89IN119-7	ZamB2 (c)

All samples analyzed at the NSSL, Lincoln, NE, unless footnoted b (Purdue University Soil Characterization Lab).

- a) Official Soil Series Type Location
- c) Map unit representative pedon
- d) Taxadjunct
- e) Soil Mechanical Lab Data

Notes to accompany the classification and correlation of the soils of Owen County, Indiana, by Byron Nagel.

Adyeville series

The Adyeville series is established by this correlation. The OSD type location is in Perry County, Indiana. Adyeville soils were included with Wellston and Muskingum soils in the 1964 soil survey.

Alvin series

The Alvin soils in the AfsB, AfsC, and AftD map units have thin strata of silt in the C horizon. The Alvin Official Soil Series description (OSD) range of characteristics (RIC) will not be changed based on data from this subset. No changes will be made to the component tables.

Arney series

The Arney series is established by this correlation. The OSD type location is the former OSD type location for the Muren series (Clay Co., IN). The Muren OSD type location is to be moved to a site that is representative of the series concept.

Belknap series

The Belknap soils (characterization data on 1 pedon) are indicated to have a regular decrease in organic carbon content. They classify into the Endo great group class. This difference does not significantly affect the use and management of these soils. They are considered taxadjuncts.

Bloomfield series

The Bloomfield soils (characterization data on 1 pedon) are indicated to be in the coarse-loamy particle-size family by about 2 percent. This is considered within allowable lab error and this soil is not considered a taxadjunct.

Bonnie series

The pH of the subhorizon immediately below the surface layer in farmed areas ranges to neutral because of the application of lime. This difference does not significantly affect the use and management of these soils. They are not considered taxadjuncts, and no changes will be made in the component table.

Chetwynd series

The Chetwynd soils in the HesG map unit have a pH range of neutral in the series control section. They are considered to have a base status above 35 percent, and therefore within the Alfisol order. This difference does not significantly affect the use and management of these soils. They are not considered taxadjuncts. The pH range is adjusted in the component table.

Cincinnati series

The Cincinnati soils in map units CkkD2 and CkkD3 have a subhorizon that meets the criteria for a fragipan, except for the bleached vertical streaks average less than 10 cm apart. This difference does not significantly affect the use and management of these soils. They are considered taxadjuncts.

Corydon series

The Corydon OSD type location is moved to a more representative area in Owen county.

Dubois series

The Dubois soils have a subhorizon that meets the criteria for a fragipan, except for the bleached vertical streaks average less than 10 cm apart. This difference does not significantly affect the use and management of these soils. They are considered taxadjuncts.

Gallimore series

The Gallimore series is established by this correlation. Gallimore soils were included with Negley soils in the 1964 soil survey. The classification of this series is not clearly defined. The OSD characterization data indicates this soil to be a Typic Dystrochrept. A pedon sampled from Morgan Co. indicates the classification to be a Typic Hapludult.

Grayford series

The Grayford soils (characterization data on 1 pedon) are indicated to have a base status above 60 percent, and therefore would classify in the typic rather than ultic subgroup. The loess thickness on the Grayford soils in map units GmcD3 and GmhD2 typically ranges from 22 to 28 inches in thickness which would classify these soils into the fine-silty particle-size family. The particle-size family of the Grayford OSD is on the break between the fine-loamy and fine-silty families. These differences do not significantly affect the use and management of these soils. They are considered taxadjuncts.

Greybrook series

The Greybrook series is established by this correlation. Greybrook soils were included with Otwell soils in the 1964 soil survey.

Haggatt series

The Haggatt series is established by this correlation. The OSD type location is in Washington County, IN. Haggatt soils are established for soils that have been correlated as Haggerstown series, bedrock substratum phase. Haggatt soils were included with Bewleyville soils in the 1964 soil survey.

Haubstadt series

The Haubstadt soils have a subhorizon that meets the criteria for a fragipan, except for the bleached vertical streaks average less than 10 cm apart. This difference does not significantly affect the use and management of these soils. They are considered taxadjuncts.

Hickory series

The Hickory soils in the HelD2, HeoE, HeuE and HeuF are indicated to have a base status less than 60 percent, and therefore would be classified in the ultic subgroup. Characterization data on 1 pedon, and 3 samples by field kit were evaluated. These soils are not considered taxadjuncts. The rock fragment content of the lower part of the series control section of the HelD2 map unit is about 5 percent more than the series range, and adjustments to the component table are made. The Hickory soils in the HefG, HesG and SwhG map units have the depth to the base of the argillic horizon and depth to carbonates at about 35 inches. These differences do not significantly affect the use and management of these soils.

Hollybrook series

The Hollybrook series is established by this correlation. Hollybrook soils were included with map units that have been strip mined and reclaimed since the 1964 soil survey.

Lieber series

The Lieber series is established by this correlation. The OSD type location is in Clay County, IN. Lieber soils were included with Vigo soils in the 1964 soil survey.

McAdoo series

The McAdoo series is established by this correlation. The OSD type location is in Posey County, IN. McAdoo soils are established for soils that have been correlated as Nolin soils that do not have a water table above 6 feet. The McAdoo soils in the MhhAK map unit do not have carbonates in the particle-size control section. This difference does not affect the use and management of this soil and it is not considered a taxadjunct. McAdoo soils were included with Genesee soils in the 1964 soil survey.

Minnehaha series

The Minnehaha series is established by this correlation. Minnehaha soils were included with strip mines in the 1964 soil survey.

Nawakwa series

The Nawakwa series is established by this correlation. Nawakwa soils were included with map units that have been strip mined and reclaimed since the 1964 soil survey.

Newark series

The Newark soils (characterization on 1 pedon) are indicated to have a regular decrease in organic carbon content to a level below 0.2 percent at a depth of 125 cm. Therefore they are considered to have a cambic horizon. This difference does not affect the use and management of this soil. They are considered taxadjuncts.

Olephant series

The Olephant series are established by this correlation. Olephant soils were included with Otwell soils in the 1964 soil survey.

Otwell series

The Otwell soils have a subhorizon that meets the criteria for a fragipan, except for the bleached vertical streaks average less than 10 cm apart. This difference does not significantly affect the use and management of these soils. They are considered taxadjuncts.

Parke series

The Parke OSD type location is moved to a more representative area in Owen County.

Patricksburg series

The Patricksburg series is established by this correlation. Patricksburg soils were included with Johnsburg soils in the 1964 soil survey.

Pekin series

The Pekin soils in the PcrAQ map unit have a subhorizon that meets the criteria for a fragipan, except for the bleached vertical streaks average less than 10 cm apart. This difference does not significantly affect the use and management of these soils. They are considered taxadjuncts.

Pike series

The Pike OSD type location is moved to a more representative area in Owen County.

Potawatomi series

The Potawatomi series is established by this correlation. Potawatomi soils were included with Tilsit soils in the 1964 soil survey.

Pottersville series

The Pottersville series is established by this correlation. Pottersville soils were included with Markland and Otwell soils in the 1964 soil survey.

Romona series

The Romona series is established by this correlation. Romona soils are included with Corydon soils in the 1964 soil survey.

Solsberry series

The Solsberry series is established by this correlation. The Solsberry soils in the SneD2, SneD3 and SneD5 map units have a subhorizon that meets the criteria for a fragipan, except for the bleached vertical streaks average less than 10 cm apart. This difference does not significantly affect the use and management of these soils. Solsberry soils were included with the Cincinnati and Zanesville soils in the 1964 soil survey. These soils are tentatively classified as Fragiudalfs, but will be evaluated in future updates and may be reclassified as Fragic Oxyaquic Hapludalfs.

Steff series

The Steff soils average less than 18 percent clay in the particle-size control section, and therefore are in the coarse-silty family. Also the upper part of the particle-size control section is non-acid in areas where lime has been applied. These differences do not significantly affect the use and management of these soils. Steff soils are considered taxadjuncts. Adjustments are made to the component table.

Stinesville series

The Stinesville series is established by this correlation. Stinesville soils were included with Grayford soils in the 1964 soil survey.

Stubenville series

The Stubenville series is established by this correlation. Stubenville soils were included with Otwell soils in the 1964 soil survey.

Taggart series

The Taggart OSD type location is moved to Morgan County from Owen County.

Tapawingo series

The Tapawingo series is established by this correlation. Tapawingo soils were included with map units that have been strip mined and reclaimed since the 1964 soil survey.

Tipsaw series

The Tipsaw series is established by this correlation. The Tipsaw OSD type location is in Perry County, IN. Tipsaw soils were included with Muskingum soils in the 1964 soil survey.

Tulip series

The Tulip series is established by this correlation. The Tulip soils were included with Muskingum and Wellston soils in the 1964 soil survey.

Vigo series

The Vigo OSD type location is moved to a more representative area in Owen County.

Wea series

The Wea soils (characterization on 1 pedon) indicate the organic-carbon content is less than 0.6 percent throughout the thickness of the mollic epipedon. This difference does not significantly affect the use and management of these soils, and they are not considered taxadjuncts.

Wellston series

The Wellston soils in the HeuE and HeuF map units are considered to average more than 15 percent fine and coarser sand in the particle-size control section. Therefore, they are in the fine-loamy family class. The higher sand content is due to the influence of glacial till above these soils. This difference does not significantly affect the use and management of these soils. They are considered taxadjuncts. Adjustments are made to the component tables.

Wilhite series

The Wilhite soils in the WolAI map unit have a regular decrease in organic carbon and a value of 0.18 at a depth of 125 cm. Therefore, this soil is indicated to have a cambic horizon and classify as a Endoaquept. This difference does not significantly affect the use and management of these soils. They are considered taxadjuncts.

Wirt series

The Wirt soils in the WpuAK map unit are slightly alkaline in the lower part of the series control section. The component table is adjusted. The Wirt soils in the WprAQ map unit have a buried soil in the lower part of the series control section. The Wirt soils in the WpyBJ map unit have a thicker surface layer than is allowed in the OSD range. The differences noted in the above map units do not significantly affect the use and management of these soils.

CLASSIFICATION OF THE SOILS

An asterick in the first column indicates that the soil is a taxadjunct to the series. See "Notes to accompany the classification and correlation of the soils of Owen County, Indiana" for a description of those characteristics of the soil that are outside the range of the series. The classification of the soils incorporates all amendments published in National Soil Taxonomy Handbook up to and including issue 17.

Soil name	Family or higher taxonomic class
.	
Adrian muck	Sandy or sandy-skeletal, mixed, euic, mesic Terric Medisaprists
Adyeville	Coarse-loamy, mixed, mesic Typic Hapludults
Alvin	Coarse-loamy, mixed, mesic Typic Hapludalfs
Armiesburg	Fine-silty, mixed, mesic Fluventic Hapludolls
Arney	Fine-silty, mixed, mesic Aquultic Hapludalfs
Ava	Fine-silty, mixed, mesic Oxyaquic Fragiudalfs
Ayrshire	Fine-loamy, mixed, mesic Aeric Endoaqualfs
*Belknap	Coarse-silty, mixed, acid, mesic Aeric Fluvaquents
Birds	Fine-silty, mixed, nonacid, mesic Typic Fluvaquents
Bloomfield	Sandy, mixed, mesic Psammentic Hapludalfs
Bonnie	Fine-silty, mixed, acid, mesic Typic Fluvaquents
Caneyville	Fine, mixed, mesic Typic Hapludalfs
Chetwynd	Fine-loamy, mixed, mesic Typic Hapludults
*Cincinnati	Fine-silty, mixed, mesic Oxyaquic Fragiudalfs
Corydon	Clayey, mixed, mesic Lithic Argiudolls
Crider	Fine-silty, mixed, mesic Typic Paleudalfs
*Dubois	Fine-silty, mixed, mesic Aeric Fragiaqualfs
Ebal	Fine, mixed, mesic Oxyaquic Hapludalfs
Elston	Coarse-loamy, mixed, mesic Typic Argiudolls
Gallimore	Coarse-loamy, mixed, mesic Typic Dystrochrepts
Gessie	Fine-loamy, mixed, mesic Fluventic Eutrochrepts
*Grayford	Fine-loamy, mixed, mesic Ultic Hapludalfs
Greybrook	Fine-loamy, mixed, mesic Typic Hapludalfs
Haggatt	Fine, mixed, mesic Typic Hapludalfs
*Haubstadt	Fine-silty, mixed, mesic Aquic Fragiudalfs
Haymond	Coarse-silty, mixed, mesic Fluventic Dystrochrepts
Hickory	Fine-loamy, mixed, mesic Typic Hapludalfs
Hollybrook	Fine-loamy, mixed, mesic Alfic Udarents
Holton	Coarse-loamy, mixed, nonacid, mesic Aeric Endoaquepts
Hoosierville	Fine-silty, mixed, mesic Typic Epiaqualfs
Lieber	Fine-silty, mixed, mesic Aeric Epiaqualfs
Martinsville	Fine-loamy, mixed, mesic Typic Hapludalfs
McAdoo	Fine-silty, mixed, mesic Fluventic Eutrochrepts
McGary	
Minnehaha	
Montgomery	Fine, mixed, mesic Vertic Endoaquolls
Moundhaven	Sandy, mixed, mesic Typic Udifluvents
Nawakwa	

SOIL SURVEY OWEN COUNTY, INDIANA

CLASSIFICATION OF THE SOILS -- Continued

Soil name	Family or higher taxonomic class
*Newark	Fine-silty, mixed, nonacid, mesic Aeric Fluvaquents
Oldenburg	Coarse-loamy, mixed, mesic Fluvaquentic Eutrochrepts
Olephant	Fine-silty, mixed, mesic Ultic Hapludalfs
*Otwell	Fine-silty, mixed, mesic Oxyaquic Fragiudalfs
Parke	Fine-silty, mixed, mesic Ultic Hapludalfs
Patricksburg	Fine-silty, mixed, mesic Aeric Fragiaqualfs
*Pekin	Fine-silty, mixed, mesic Aquic Fragiudults
Peoga	Fine-silty, mixed, mesic Typic Epiaqualfs
Piankeshaw	Fine-loamy, mixed, mesic Dystric Fluventic Eutrochrepts
Pike	Fine-silty, mixed, mesic Ultic Hapludalfs
Pope	Coarse-loamy, mixed, mesic Fluventic Dystrochrepts
Potawatomi	Fine-silty, mixed, mesic Aquic Fragiudalfs
Pottersville	Fine-silty, mixed, mesic Typic Hapludalfs
Princeton	Fine-loamy, mixed, mesic Typic Hapludalfs
Romona	Fine, mixed, mesic Typic Argiudolls
Ryker	Fine-silty, mixed, mesic Typic Paleudalfs
Shakamak	Fine-silty, mixed, mesic Aquic Fragiudalfs
Shircliff	Fine, mixed, mesic Oxyaquic Hapludalfs
Solsberry	Fine-silty, mixed, mesic Oxyaquic Fragiudalfs
*Steff	Fine-silty, mixed, mesic Fluvaquentic Dystrochrepts
Stendal	Fine-silty, mixed, acid, mesic Aeric Fluvaquents
Stinesville	Fine-silty, mixed, mesic Typic Paleudalfs
Stonelick	Coarse-loamy, mixed, calcareous, mesic Typic Udifluvents
Stubenville	Fine-silty, mixed, mesic Typic Hapludalfs
Taggart	Fine-silty, mixed, mesic Aeric Epiaqualfs
Tapawingo	Loamy, mixed, mesic Typic Udorthents
Tipsaw	
Tulip	
Udorthents	Loamy, mixed, mesic Udorthents
Vigo	
Wea	Fine-loamy, mixed, mesic Typic Argiudolls
*Wellston	Fine-silty, mixed, mesic Ultic Hapludalfs
Whitaker	Fine-loamy, mixed, mesic Aeric Endoaqualfs
Wilbur	Coarse-silty, mixed, mesic Fluvaquentic Eutrochrepts
*Wilhite	Fine, mixed, nonacid, mesic Typic Fluvaquents
Wirt	Coarse-loamy, mixed, mesic Dystric Fluventic Eutrochrepts
Zanesville	Fine-silty, mixed, mesic Oxyaquic Fragiudalfs

Certifications

The State Soil Scientist certifies that:

- a) The field mapping was completed in October 1993.
- b) Interpretations have been coordinated with adjoining survey areas.
- c) The location of all typical pedons in the survey area are correct and are within delineation's that have the referenced name.
- d) All typical pedons are correctly classified according to Soil Taxonomy, and amendment issues 1-17 of the National Soil Taxonomy Handbook.
- e) The soil maps are complete, accurate and consistent.
- f) Owen County has made a quality join with the following soil survey areas: Clay, Greene, Monroe, Morgan and Putnam counties. The Correlation Memorandums for these survey areas will not be amended at this time. A record of the map unit changes is placed on a set of the soil maps for each county. These revised maps will be filed in the MLRA Project Office in each county case file.

Clay County (published 1982); The Clay County Soil Survey will accept the following Owen Co. map units:

AfsB	Alvin-Princeton fine sandy loams, 2 to 6 percent slopes
AfsC	Alvin-Princeton fine sandy loams, 6 to 12 percent slopes
BdxAH	Belknap silt loam, 0 to 2 percent slopes, frequently flooded, very brief duration $% \left(1\right) =\left(1\right) +\left(1\right) +$
CkkD2	Cincinnati silt loam, 12 to 18 percent slopes, eroded
CkkD3	Cincinnati silt loam, 12 to 18 percent slopes, severely eroded
GabG	Gallimore-Chetwynd loams, 25 to 70 percent slopes
HccB2	Haubstadt silt loam, 2 to 6 percent slopes, eroded
HeoE	Hickory silt loam, 18 to 25 percent slopes
HeoG	Hickory silt loam, 25 to 70 percent slopes
HepG	Hickory-Adyeville complex, 35 to 60 percent slopes
HeuE	Hickory-Wellston silt loams, 18 to 25 percent slopes
HeuF	Hickory-Wellston silt loams, 25 to 35 percent slopes
HleAH	Holton silt loam, 0 to 2 percent slopes, frequently flooded, very brief duration $% \left(1\right) =\left(1\right) +\left(
MhhAK	McAdoo silt loam, 0 to 2 percent slopes, occasionally flooded, brief duration

MrcG	Minnehaha silty clay loam, 35 to 70 percent slopes
NbhAH	Newark silt loam, 0 to 2 percent slopes, frequently flooded, brief duration $\ \ $
OfaAH	Oldenburg silt loam, 0 to 2 percent slopes, frequently flooded, very brief duration
OmkC3	Otwell silt loam, 6 to 12 percent slopes, severely eroded
PbbC2	Parke silt loam, 6 to 12 percent slopes, eroded
PlfB2	Pike silt loam, 2 to 6 percent slopes, eroded
SfoA	Shakamak silt loam, 1 to 3 percent slopes
SneC2	Solsberry silt loam, 6 to 12 percent slopes, eroded
SneC3	Solsberry silt loam, 6 to 12 percent slopes, severely eroded
SneD2	Solsberry silt loam, 12 to 18 percent slopes, eroded
SneD3	Solsberry silt loam, 12 to 18 percent slopes, severely eroded
StaAH	Steff silt loam, 0 to 2 percent slopes, frequently flooded, very brief duration
StdAH	Stendal silt loam, 0 to 2 percent slopes, frequently flood, brief duration
TtcE	Tulip-Wellston-Adyeville silt loams, 18 to 25 percent slopes
WokAH	Wilbur silt loam, 0 to 2 percent slopes, frequently flooded, brief duration $% \left(1\right) =\left(1\right) +\left(1\right)$
WpuAH	Wirt silt loam, 0 to 2 percent slopes, frequently flooded, very brief duration $\ \ $

Greene county (published 1988); the Greene county Soil Survey will accept the following Owen co. map units:

AffE	Alvin-Bloomfield-Princeton complex, 18 to 25 percent slopes
AfsB	Alvin-Princeton fine sandy loams, 2 to 6 percent slopes
AfsC	Alvin-Princeton fine sandy loams, 6 to 12 percent slopes
GaaE2	Gallimore loam, 18 to 25 percent slopes, eroded
GabG	Gallimore-Chetwynd loams, 25 to 70 percent slopes
GmpE2	Greybrook silt loam, 18 to 25 percent slopes, eroded
HasE2	Haggatt-Caneyville silt loams, 18 to 25 percent slopes, eroded
HeoG	Hickory silt loam, 25 to 70 percent slopes
HepG	Hickory-Adyeville complex, 35 to 60 percent slopes
HeuE	Hickory-Wellston silt loams, 18 to 25 percent slopes

HeuF Hickory-Wellston silt loams, 25 to 35 percent slopes MhhAH McAdoo silt loam, 0 to 2 percent slopes, frequently flooded, brief duration NbhAH Newark silt loam, 0 to 2 percent slopes, frequently flooded, brief duration PcrC2 Pekin silt loam, 6 to 12 percent slopes, eroded Pml Pits, quarries Potawatomi silt loam, 1 to 3 percent slopes PryB Pottersville silt loam, 12 to 18 percent slopes, severely eroded PsaD3 Solsberry silt loam, 6 to 12 percent slopes, eroded SneC2 SneC3 Solsberry silt loam, 6 to 12 percent slopes, severely eroded SneD2 Solsberry silt loam, 12 to 18 percent slopes, eroded SneD3 Solsberry silt loam, 12 to 18 percent slopes, severely eroded StdAH Stendal silt loam, 0 to 2 percent slopes, frequently flooded, brief duration Stonelick sandy loam, 0 to 2 percent slopes, frequently flooded, SupAH brief duration TcgG Tipsaw-Rock outcrop complex, 35 to 70 percent slopes Tulip-Tipsaw complex, 25 to 60 percent slopes TtaG TtcE Tulip-Wellston-Adyeville silt loams, 18 to 25 percent slopes Wilhite silty clay loam, 0 to 1 percent slopes, frequently WomAH flooded, brief duration Wirt silt loam, 0 to 2 percent slopes, frequently flooded, very WpuAH brief duration ZamB2 Zanesville silt loam, soft bedrock substratum, 2 to 6 percent slopes, eroded Zanesville silt loam, soft bedrock substratum, 6 to 12 percent ZamC2 slopes, eroded ZamC3 Zanesville silt loam, soft bedrock substratum, 6 to 12 percent slopes, severely eroded Zanesville, soft bedrock substratum-Tulip silt loams, 12 to 18 ZapD3 percent slopes, severely eroded

Monroe County (published 1981); the Monroe County Soil Survey will accept the following Owen County map units:

BdxAH Belknap silt loam, 0 to 2 percent slopes, frequently flooded, very brief duration

CspC2 Crider silt loam, 6 to 12 percent slopes, eroded Crider silt loam, 6 to 12 percent slopes, severely eroded CspC3 Dubois silt loam, 0 to 2 percent slopes DfnA GabG Gallimore-Chetwynd loams, 25 to 70 percent slopes HarD2 Haggatt silt loam, 12 to 18 percent slopes, eroded HarD3 Haggatt silt loam, 12 to 18 percent slopes, severely eroded HasE2 Haggatt-Caneyville silt loams, 18 to 25 percent slopes, eroded HccB2 Haubstadt silt loam, 2 to 6 percent slopes, eroded HcgAH Haymond silt loam, 0 to 2 percent slopes, frequently flooded, brief duration HeoG Hickory silt loam, 25 to 70 percent slopes LnfA Lieber silt loam, 0 to 2 percent slopes MsvA Montgomery silty clay loam, 0 to 1 percent slopes NbhAH Newark silt loam, 0 to 2 percent slopes, frequently flooded, brief duration OmkC2 Otwell silt loam, 6 to 12 percent slopes, eroded PcrC2 Pekin silt loam, 6 to 12 percent slopes, eroded PryB Potawatomi silt loam, 1 to 3 percent slopes St.fB2 Stinesville silt loam, 2 to 6 percent slopes, eroded StfC2 Stinesville silt loam, 6 to 12 percent slopes, eroded SupAH Stonelick sandy loam, 0 to 2 percent slopes, frequently flooded, brief duration TtaG Tulip-Tipsaw complex, 25 to 60 percent slopes Tulip-Wel1ston-Adyeville silt loams, 18 to 25 percent TtcE slopes WhfD2 Wellston silt loam, 12 to 18 percent slopes, eroded WokAH Wilbur silt loam, 0 to 2 percent slopes, frequently flooded, brief duration WpuAH Wirt silt loam, 0 to 2 percent slopes, frequently flooded, very brief duration Zanesville silt loam, soft bedrock substratum, 6 to 12 percent ZamC2 slopes, eroded Zanesville silt loam, soft bedrock substratum, 6 to 12 percent ZamC3 slopes, severely eroded

Morgan County (published 1981); the Morgan County Soil Survey will accept the following Owen Co. map units:

AhrAK Armiesburg silty clay loam, 0 to 2 percent slopes, occasionally flooded, brief duration

AloB2 Ava silt loam, 2 to 6 percent slopes, eroded

DfnA Dubois silt loam, 0 to 2 percent slopes

GabG Gallimore-Chetwynd loams, 25 to 70 percent slopes

GmhD2 Grayford-Ryker silt loams, 12 to 18 percent slopes, eroded

HccB2 Haubstadt silt loam, 2 to 6 percent slopes, eroded

HchAP Haymond silt loam, sinkhole, 0 to 2 percent slopes, occasionally

flooded, very brief duration

HeoG Hickory silt loam, 25 to 70 percent slopes

LnfA Lieber silt loam, 0 to 2 percent slopes

NbhAH Newark silt loam, 0 to 2 percent slopes, frequently flooded,

brief duration

OmkC2 Otwell silt loam, 6 to 12 percent slopes, eroded

PhaA Peoga silt loam, 0 to 1 percent slopes

PlcAH Piankeshaw silt loam, 0 to 2 percent slopes, frequently

flooded, very brief duration

PsaG Pottersville silt loam, 25 to 50 percent slopes

RpzG Romona-Corydon-Rock outcrop complex, 35 to 60 percent slopes

SfyB2 Shircliff silt loam, 2 to 6 percent slopes, eroded

StfB2 Stinesville silt loam, 2 to 6 percent slopes, eroded

StfC2 Stinesville silt loam, 6 to 12 percent slopes, eroded

StgC2 Stinesville-Ryker-Grayford silt loams, karst, rolling, eroded

StgD2 Stinesville-Ryker-Grayford silt loams, karst, hilly, eroded

TcgG Tipsaw-Rock outcrop complex, 35 to 70 percent slopes

WeaAQ Wea, sandy substratum-Elston complex, 0 to 2 percent slopes,

rarely flooded, very brief duration

WlmAK Whitaker loam, 0 to 2 percent slopes, occasionally flooded,

brief duration

WokAH Wilbur silt loam, 0 to 2 percent slopes, frequently flooded, brief

duration

Putnam county (published 1981); the Putnam County Soil Survey will accept the following Owen County map units:

AkeB2 Arney silt loam, 2 to 6 percent slopes, eroded

AloB2 Ava silt loam, 2 to 6 percent slopes, eroded

BgeAH Birds silt loam, 0 to 1 percent slopes, frequently flooded, brief

duration

DfnA Dubois silt loam, 0 to 2 percent slopes

GmhD2 Grayford-Ryker silt loams, 12 to 18 percent slopes, eroded

GmpF Greybrook silt loam, 25 to 35 percent slopes

HccB2 Haubstadt silt loam, 2 to 6 percent slopes, eroded

HcgAH Haymond silt loam, 0 to 2 percent slopes, frequently flooded,

brief duration

HefG Hickory loam, 45 to 70 percent slopes

HeoG Hickory silt loam, 25 to 70 percent slopes

HesG Hickory-Chetwynd loams, 35 to 70 percent slopes

HeuE Hickory-Wellston silt loams, 18 to 25 percent slopes

HeuF Hickory-Wellston silt loams, 25 to 35 percent slopes

HleAH Holton silt loam, 0 to 2 percent slopes, frequently flooded,

very brief duration

LnfA Lieber silt loam, 0 to 2 percent slopes

NbhAH Newark silt loam, 0 to 2 percent slopes, frequently flooded,

brief duration

OfeB Olephant silt loam, 1 to 3 percent slopes

OmkC2 Otwell silt loam, 6 to 12 percent slopes, eroded

OmkC3 Otwell silt loam, 6 to 12 percent slopes, severely eroded

OmkD3 Otwell silt loam, 12 to 18 percent slopes, severely eroded

PhaA Peoga silt loam, 0 to 1 percent slopes

PlfB2 Pike silt loam, 2 to 6 percent slopes, eroded

StfB2 Stinesville silt loam, 2 to 6 percent slopes, eroded

StfC2 Stinesville silt loam, 6 to 12 percent slopes, eroded

WpyBJ Wirt-Pekin silt loams, 0 to 6 percent slopes, frequently

flooded, very long duration

General Soil Map

A 1:250,000 STATSGO map was used as the base map for the general soil map. This map will be used to update all adjoining subsets. Therefore, a general soil map join was not made with the adjoining subsets.

Approval Signature and Date

TRAVIS NEELY / Pate
Soil Survey Area 11 Team Leader

ROBERT L. EDDLEMAN / Date

Edutt Eddleman 5/0/17

State Conservationist